

# Towards Homotopical Algebraic Quantum Field Theory

Alexander Schenkel

School of Mathematical Sciences, University of Nottingham



The University of  
**Nottingham**

UNITED KINGDOM · CHINA · MALAYSIA



THE ROYAL SOCIETY

Talk @ Foundational and Structural Aspects of Gauge Theories,  
Mainz Institute for Theoretical Physics, 29 May – 2 June 2017.

Based on works with different subsets of

Collaborators := {C. Becker, M. Benini, U. Schreiber, R. J. Szabo}

1. Explain why

AQFT/LCQFT is insufficient to describe gauge theories

2. Present ideas/observations indicating that the key to resolve this problem is

homotopical LCQFT := homotopical algebra + LCQFT

3. Discuss our results and inform you about the state-of-the-art of our development of homotopical LCQFT

# LCQFT vs Gauge Theory

# LCQFT = AQFT on Lorentzian manifolds

- ◇ **Basic idea** [Brunetti, Fedenhagen, Verch; ...]


$$\begin{array}{ccc} \text{Loc} & \xrightarrow{\text{functor } \mathfrak{A}} & \text{Alg} \\ \text{category of spacetimes} & & \text{category of algebras} \end{array}$$

↪ “Coherent assignment of observable algebras to spacetimes”

- $\mathfrak{A}(M)$  = observables we can measure in  $M$
- $\mathfrak{A}(f) : \mathfrak{A}(M) \rightarrow \mathfrak{A}(M')$  = embedding of observables along  $f : M \rightarrow M'$


- ◇ **BFV axioms** (motivated from physics)

Isotony:




$$\mathfrak{A}(M) \xrightarrow{\text{mono}} \mathfrak{A}(M')$$

Causality:



$$[\mathfrak{A}(M_1), \mathfrak{A}(M_2)] = \{0\}$$

Time-slice:



$$\mathfrak{A}(M) \xrightarrow{\text{iso}} \mathfrak{A}(M')$$

# Local-to-global property

For every spacetime  $M$ , the global algebra  $\mathfrak{A}(M)$  can be “recovered” from the algebras  $\mathfrak{A}(U)$  corresponding to suitable sub-spacetimes  $U \subseteq M$ .

◇ Different ways to formalize this property:

1. **Cosheaf property:**  $\mathfrak{A} : \text{Loc} \rightarrow \text{Alg}$  is cosheaf (w.r.t. suitable topology)

⚡ only true for extremely special covers  $\Rightarrow$  too strong condition


2. **Additivity:**  $\mathfrak{A}(M) \simeq \bigvee_{\alpha} \mathfrak{A}(U_{\alpha})$  for suitable covers  $\{U_{\alpha} \subseteq M\}$  [Fewster; ...]

✓ true in examples ⚡ need to know  $\mathfrak{A}(M)$

3. **Universality:**  $\mathfrak{A}(M)$  is isomorphic to *Fredenhagen's universal algebra* corresponding to  $\{U \subseteq M : \text{open, causally compatible and } U \simeq \mathbb{R}^m\}$

✓  $\mathfrak{A}$  determined by restriction  $\mathfrak{A}_{\odot} : \text{Loc}_{\odot} \rightarrow \text{Alg}$  via left Kan extension

✓ true in examples [Lang]



# Does $U(1)$ -Yang-Mills theory fit into LCQFT?

- ◇ **Differential cohomology groups** = gauge orbit spaces

$$\widehat{H}^2(M) \cong \frac{\{ \text{principal } U(1)\text{-bundles } P \rightarrow M \text{ with connection } A \}}{\{ \text{gauge transformations} \}}$$

- ◇ Solution spaces of  $U(1)$ -Yang-Mills theory

$$\mathcal{F}(M) := \{ h \in \widehat{H}^2(M) : \delta \text{curv}(h) = 0 \}$$

are Abelian Fréchet-Lie groups with natural presymplectic structure  $\omega_M$


## Theorem [Becker,AS,Szabo:1406.1514]

Quantization of smooth Pontryagin dual of  $(\mathcal{F}(M), \omega_M)$  defines functor  $\mathfrak{Q} : \text{Loc} \rightarrow \text{Alg}$  which satisfies **causality** and **time-slice**, but violates **isotony** and **local-to-global properties**.


**NB:** Similar results for  $S$ -duality invariant theory [Becker,Benini,AS,Szabo:1511.00316] and also for less complete approaches based on  $A$ -fields or  $F$ -fields [Sanders,Dappiaggi,Hack; Fewster,Lang; ...]

# What is the source of these problems?

1. Isotony fails because gauge theories carry **topological charges**



2. Local-to-global property fails because we took **gauge invariant observables**



1. Violation of isotony is a physical feature, hence we have to accept that!
2. Violation of local-to-global property is an artifact of our description by gauge invariant observables, hence we can improve that!


# Groupoids vs Gauge Orbit Spaces



# Groupoids of gauge fields

- ◇ Let's consider for the moment gauge theory on  $M \simeq \mathbb{R}^m$ 
  1. **Gauge fields**  $A \in \Omega^1(M, \mathfrak{g})$
  2. **Gauge transformations**  $g \in C^\infty(M, G)$  acting as  $A \triangleleft g = g^{-1} A g + g^{-1} dg$
- ◇ **Groupoid** of gauge fields on  $M$

$$\mathcal{G}(M) := \Omega^1(M, \mathfrak{g}) \rtimes C^\infty(M, G) =$$



Two groupoids are “the same” not only when isomorphic, but also when **weakly equivalent**  $\rightsquigarrow$  **model category/homotopical algebra**

- ◇ Non-redundant information encoded in the groupoid  $\mathcal{G}(M)$ 
  1. Gauge orbit space  $\pi_0(\mathcal{G}(M)) = \Omega^1(M, \mathfrak{g})/C^\infty(M, G)$
  2. Automorphism groups  $\pi_1(\mathcal{G}(M), A) = \{g \in C^\infty(M, G) : A \triangleleft g = A\}$

**!** Gauge invariant observables ignore the  $\pi_1$ 's, hence are incomplete!

# Groupoids and local-to-global properties

- ◇ Groupoids of gauge fields satisfy very strong local-to-global property

## Homotopy sheaf property

For all manifolds  $M$  and all open covers  $\{U_\alpha \subseteq M\}$ , the canonical map

$$\mathcal{G}(M) \longrightarrow \operatorname{holim} \left( \prod_{\alpha} \mathcal{G}(U_{\alpha}) \rightrightarrows \prod_{\alpha\beta} \mathcal{G}(U_{\alpha\beta}) \Rrightarrow \prod_{\alpha\beta\gamma} \mathcal{G}(U_{\alpha\beta\gamma}) \Rrightarrow \cdots \right)$$

is a weak equivalence in  $\operatorname{Grpd}$ .

- ◇ Precise formulation of the familiar “gluing up to gauge transformation”

$$\left\{ (\{A_{\alpha}\}, \{g_{\alpha\beta}\}) : A_{\beta}|_{U_{\alpha\beta}} = A_{\alpha}|_{U_{\alpha\beta}} \triangleleft g_{\alpha\beta}, \quad g_{\alpha\beta} g_{\beta\gamma} = g_{\alpha\gamma} \text{ on } U_{\alpha\beta\gamma} \right\}$$
$$\Downarrow 1:1$$
$$\left\{ \text{gauge fields on } M \right\}$$

- ◇ **Crucial Point:** Taking into account the **groupoids** of gauge fields, rather than only the **gauge orbit spaces**, there are very strong homotopical local-to-global properties for classical gauge theories!

# Cosimplicial observable algebras

# What are “function algebras” on groupoids?

- ◇ QFT needs quantized ‘algebras’ of functions on the ‘spaces’ of fields

- ✓ Space of fields  $\mathcal{F}(M)$  is set (+ smooth structure)

- $\rightsquigarrow \mathcal{O}(M) = C^\infty(\mathcal{F}(M))$  has the structure of an algebra

- ? Space of fields  $\mathcal{G}(M)$  is groupoid (+ smooth structure)

- $\rightsquigarrow \mathcal{O}(M) = “C^\infty(\mathcal{G}(M))” = ?$  has which algebraic structure?

- ◇ Nerve construction  $N : \text{Grpd} \rightarrow \text{sSet}$  assigns the simplicial set

$$N(\mathcal{G}(M)) = \left( \Omega^1(M, \mathfrak{g}) \rightrightarrows \Omega^1(M, \mathfrak{g}) \times C^\infty(M, G) \rightrightarrows \dots \right)$$

- ◇ Taking level-wise smooth functions gives **cosimplicial algebra**

$$\mathcal{O}(M) = \left( C^\infty(\Omega^1(M, \mathfrak{g})) \rightrightarrows C^\infty(\Omega^1(M, \mathfrak{g}) \times C^\infty(M, G)) \rightrightarrows \dots \right)$$

# Relation to the BRST formalism and ghost fields

- ◇ Dual Dold-Kan correspondence gives equivalence  $\text{cAlg} \rightleftarrows \text{dgAlg}^{\geq 0}$
- ⇒ Equivalent description of  $\mathcal{O}(M)$  in terms of **differential graded algebra**

$$\mathcal{O}_{\text{dg}}(M) = \left( C^\infty(\Omega^1(M, \mathfrak{g})) \xrightarrow{d} C^\infty(\Omega^1(M, \mathfrak{g}) \times C^\infty(M, G)) \xrightarrow{d} \dots \right)$$

- ◇ Considering only infinitesimal gauge transformations  $C^\infty(M, \mathfrak{g})$

$$\mathcal{O}_{\text{dg}}(M) \xrightarrow{\text{van Est map}} \underbrace{C^\infty(\Omega^1(M, \mathfrak{g}))}_{\text{gauge field observables}} \otimes \underbrace{\wedge^\bullet C^\infty(M, \mathfrak{g})^*}_{\text{ghost field observables}}$$

The cosimplicial algebra  $\mathcal{O}(M)$  (or equivalently our dg-algebra  $\mathcal{O}_{\text{dg}}(M)$ ) describes non-infinitesimal analogs  $C^\infty(M, G)$  of ghost fields  $C^\infty(M, \mathfrak{g})$

- ⇒ **BRST formalism for finite gauge transformations**

# Working definition for homotopical LCQFT

# Working definition (intentionally imprecise)

A **homotopical LCQFT** is a (weak) functor  $\mathfrak{A} : \text{Loc} \rightarrow \text{dgAlg}^{\geq 0}$  to the model category of noncommutative dg-algebras, which satisfies the following axioms:

1. *Causality*: Given causally disjoint  $M_1 \xrightarrow{f_1} M \xleftarrow{f_2} M_2$ , there exist a (coherent) cochain homotopy  $\lambda_{f_1, f_2}$  such that

$$[\cdot, \cdot]_{\mathfrak{A}(M)} \circ (\mathfrak{A}(f_1) \otimes \mathfrak{A}(f_2)) = \lambda_{f_1, f_2} \circ d + d \circ \lambda_{f_1, f_2}$$

2. *Time-slice*: Given Cauchy morphism  $f : M \rightarrow M'$ , there exists a (coherent) homotopy inverse  $\mathfrak{A}(f)^{-1}$  of  $\mathfrak{A}(f)$ .
3. *Universality*:  $\mathfrak{A} : \text{Loc} \rightarrow \text{dgAlg}^{\geq 0}$  is the homotopy left Kan extension of its restriction  $\mathfrak{A}_{\odot} : \text{Loc}_{\odot} \rightarrow \text{dgAlg}^{\geq 0}$ .

**Rem:** 'Coherent' in e.g. 1.) means that the homotopies for different commutations of more than 2 observables (e.g.  $abc \rightarrow acb \rightarrow cab$  vs  $abc \rightarrow cab$ ) coincide up to specified higher homotopies.

Precise definition requires **colored operads** [Benini, AS, Woike: [work in progress](#)]

homotopical LCQFT := LCQFT $_{\infty}$ -algebra + operadic universality

# Local-to-global property in Abelian gauge theory



# Universal global gauge theory observables

- ◇ For  $G = U(1)$  and  $M \simeq \mathbb{R}^m$ ,  $\mathcal{G}(M)$  can be described by chain complex

$$\mathcal{G}_{\text{chain}}(M) = \left( \Omega^1(M) \xleftarrow{\frac{1}{2\pi i} \text{d log}} C^\infty(M, U(1)) \right)$$

- ◇ Smooth Pontryagin dual cochain complex of observables

$$\mathcal{O}_{\odot}(M) := \left( \Omega_c^{m-1}(M) \xrightarrow{\text{d}} \Omega_{c;\mathbb{Z}}^m(M) \right)$$

- ◇ Homotopy left Kan extension of  $\mathcal{O}_{\odot} : \text{Loc}_{\odot} \rightarrow \text{Ch}^{\geq 0}$

$$\mathcal{O}(M) := \text{hocolim} \left( \mathcal{O}_{\odot} : \text{Loc}_{\odot} \downarrow M \rightarrow \text{Ch}^{\geq 0} \right)$$

## Theorem [Benini,AS,Szabo:1503.08839]

1. For  $M \simeq \mathbb{R}^m$ ,  $\mathcal{O}_{\odot}(M)$  and  $\mathcal{O}(M)$  are naturally weakly equivalent.
2. For every  $M$ ,  $\mathcal{O}(M)$  weakly equivalent to dual Deligne complex on  $M$ .

- ◇ **Crucial Point:** Our homotopical version of “Fredenhagen’s universal algebra” produces the correct global observables in Abelian gauge theory, in contrast to the non-homotopical version [Dappiaggi,Lang; Fewster,Lang]!

# Toy-models of homotopical LCQFT

# LCQFT on structured spacetimes

## ◇ Basic idea [Benini,AS:1610.06071]

1. Consider LCQFT  $\mathfrak{A} : \mathbf{Str} \rightarrow \mathbf{Alg}$  on category of **spacetimes with extra geometric structures**, i.e. category fibered in groupoids  $\pi : \mathbf{Str} \rightarrow \mathbf{Loc}$ . ( $\pi^{-1}(M)$  is groupoid of structures over  $M$ , e.g. spin structures, gauge fields)
2. Regard  $\mathfrak{A}$  as a trivial homotopical LCQFT  $\mathfrak{A} : \mathbf{Str} \rightarrow \mathbf{dgAlg}^{\geq 0}$  via embedding  $\mathbf{Alg} \rightarrow \mathbf{dgAlg}^{\geq 0}$  of algebras into dg-algebras.
3. Perform homotopy right Kan extension

$$\begin{array}{ccc} \mathbf{Str} & \xrightarrow{\mathfrak{A}} & \mathbf{dgAlg}^{\geq 0} \\ & \searrow \pi & \nearrow \uparrow \\ & & \mathbf{Loc} \end{array}$$

$\text{hoU}_{\pi} \mathfrak{A}$

to induce a **nontrivial** homotopical LCQFT  $\text{hoU}_{\pi} \mathfrak{A}$  on  $\mathbf{Loc}$ .

- ◇ **Physical interpretation:** Homotopy right Kan extension turns the background fields described by  $\pi^{-1}(M)$  into observables in  $\text{hoU}_{\pi} \mathfrak{A}(M)$ .

# Properties of $\mathrm{hoU}_\pi \mathfrak{A}$

- ◇ Explicit description of degree 0 of  $\mathrm{hoU}_\pi \mathfrak{A}(M)$

$$\mathrm{hoU}_\pi \mathfrak{A}(M)^0 = \prod_{S \in \pi^{-1}(M)} \mathfrak{A}(S) \ni \left( a : \pi^{-1}(M) \ni S \mapsto a(S) \in \mathfrak{A}(S) \right)$$

- ◇ **Physical interpretation:** Combination of **classical gauge field observables** and **quantum matter field observables!**

## Theorem [Benini,AS:1610.06071]

Assume that  $\pi : \mathrm{Str} \rightarrow \mathrm{Loc}$  is strongly Cauchy flabby. Then the homotopy right Kan extension  $\mathrm{hoU}_\pi \mathfrak{A} : \mathrm{Loc} \rightarrow \mathrm{dgAlg}^{\geq 0}$  satisfies the **causality and time-slice axioms of homotopical LCQFT**. (Coherences just established in low orders.)

- ✓ First toy-models satisfying the new homotopical LCQFT axioms!  
(Proving universality is hard:  $\mathrm{hocolim}$ 's in  $\mathrm{dgAlg}^{\geq 0}$  are beyond our current technology.)

# Stack of non-Abelian Yang-Mills fields

# Yang-Mills stack

- ◇ **Motivation:** Prior to deformation quantization, we have to understand the **geometry of the groupoid of Yang-Mills solutions** and the Cauchy problem
- ↪ **Stacks**  $\cong$  presheaves of groupoids  $X$  on  $\text{Cart}$  satisfying descent [Hollander]
- ◇ **Basic idea:** Smooth structure on  $X$  is encoded by specifying groupoid  $X(\mathbb{R}^k)$  of *all* smooth maps  $\mathbb{R}^k \rightarrow X$  for *all*  $\mathbb{R}^k$  in  $\text{Cart}$  (**functor of points**)
- ◇  $\exists$  abstract model categorical construction of the stack of non-Abelian Yang-Mills solutions  $G\text{Sol}(M)$  [Benini,AS,Schreiber:1704.01378]
- ◇ Explicit description of  $G\text{Sol}(M)$  up to weak equivalence

$$G\text{Sol}(M)(\mathbb{R}^k) = \begin{cases} \text{obj} : & \text{smoothly } \mathbb{R}^k\text{-parametrized Yang-Mills solutions } (\mathbf{A}, \mathbf{P}) \\ \text{mor} : & \text{smoothly } \mathbb{R}^k\text{-parametrized gauge transformations} \\ & \mathbf{h} : (\mathbf{A}, \mathbf{P}) \rightarrow (\mathbf{A}', \mathbf{P}') \end{cases}$$

- ◇ For  $M \simeq \mathbb{R}^m$  even simpler in terms of vertical geometry on  $M \times \mathbb{R}^k \rightarrow \mathbb{R}^k$   
 $(\mathbf{A}, \mathbf{P}) = A \in \Omega^{1,0}(M \times \mathbb{R}^k, \mathfrak{g}) \quad \text{s.t.} \quad \delta_A^{\text{vert}} F^{\text{vert}}(A) = 0$

# Stacky Cauchy problem

- ◇  $\exists$  map of stacks  $\text{data}_\Sigma : \text{GSol}(M) \rightarrow \text{GData}(\Sigma)$  assigning to Yang-Mills solutions their initial data on Cauchy surface  $\Sigma \subseteq M$

**Def:** The **stacky Cauchy problem** is well-posed if  $\text{data}_\Sigma$  is a weak equivalence.

## Theorem [Benini,AS,Schreiber:1704.01378]

The stacky Yang-Mills Cauchy problem is well-posed if and only if the following hold true, for all  $\mathbb{R}^k$  in Cart:

1. For all  $(\mathbf{A}^\Sigma, \mathbf{E}, \mathbf{P}^\Sigma)$  in  $\text{GData}(\Sigma)(\mathbb{R}^k)$ , there exists  $(\mathbf{A}, \mathbf{P})$  in  $\text{GSol}(M)(\mathbb{R}^k)$  and iso  $\mathbf{h}^\Sigma : \text{data}_\Sigma(\mathbf{A}, \mathbf{P}) \rightarrow (\mathbf{A}^\Sigma, \mathbf{E}, \mathbf{P}^\Sigma)$  in  $\text{GData}(\Sigma)(\mathbb{R}^k)$ .
2. For any other iso  $\mathbf{h}'^\Sigma : \text{data}_\Sigma(\mathbf{A}', \mathbf{P}') \rightarrow (\mathbf{A}^\Sigma, \mathbf{E}, \mathbf{P}^\Sigma)$  in  $\text{GData}(\Sigma)(\mathbb{R}^k)$ , there exists **unique** iso  $\mathbf{h} : (\mathbf{A}, \mathbf{P}) \rightarrow (\mathbf{A}', \mathbf{P}')$  in  $\text{GSol}(M)(\mathbb{R}^k)$ , such that  $\mathbf{h}'^\Sigma \circ \text{data}_\Sigma(\mathbf{h}) = \mathbf{h}^\Sigma$ .

! Interesting **smoothly  $\mathbb{R}^k$ -parametrized Cauchy problems!** To the best of my knowledge, positive results only known for  $\mathbb{R}^0$  [Chrusciel,Shatah; Choquet-Bruhat].

# Summary and Outlook



# Summary and Outlook

- ◇ Quantum gauge theories are **NOT** contained in the LCQFT framework
- ◇ To capture crucial homotopical features of classical gauge theories, one needs “higher algebras” to formalize quantum gauge theories
  - ⇒ **Homotopical LCQFT**
- ◇ Already very promising results:
  - ✓ Local-to-global property of observables [Benini,AS,Szabo:1503.08839]
  - ✓ Toy-models of homotopical LCQFT [Benini,AS:1610.06071]
  - ✓ Yang-Mills stack and stacky Cauchy problem [Benini,AS,Schreiber:1704.01378]
- ◇ Open problems/Work in progress:
  1. Develop operadic approach to homotopical LCQFT to control coherences
  2. Construct proper examples of dynamical and quantized gauge theories
  3. What's the physics behind “higher algebras”? [Thanks for asking, Klaus!]

**Thanks for your attention.**